

Dipole in Shell


Kirk T. McDonald

Joseph Henry Laboratories, Princeton University, Princeton, NJ 08544

(October 2, 2001)

1 Problem

What is the (vector) electric field strength at the center of a spherical cavity of radius a in a grounded conductor if a point dipole \mathbf{p} is placed at distance b ($0 < b < a$) from the center of the cavity so as to make angle α to the radius vector to the dipole?



2 Solution

This problem is question 05/01 at Yakov Kantor's Physics Quiz site,

<http://www.tau.ac.il/~kantor/QUIZ/>

We solve via the image method for spherical geometry. Recall that a charge q at radius $b < a$ results in an image charge q' at a^2/b , such that the potential at $r = a$ vanishes,


$$\phi(r = a) = 0 = \frac{q}{a - b} + \frac{q'}{a^2/b - a} = \frac{q}{a - b} + \frac{q'b/a}{a - b}, \quad (1)$$

and hence $q' = -qa/b$.

For the present case of a point dipole \mathbf{p} , we can think of this as due to charges $\pm q$ separated by distance $\delta = q/p$, with the charge $-q$ at radius b and the charge $+q$ at radius $b + \delta \cos \alpha$ as shown below.

The image transformation $r \rightarrow a^2/r$ is conformal (angle preserving), so the triangle formed by the center of the sphere and charges $\pm q$ is similar to that formed by the center and the image charges q' and q'' . Image charge $q' = +qa/b$ is at distance a^2/b from the center. Hence, the distance between the two image charges is,

$$\delta' = \delta \frac{a^2}{b^2}. \quad (2)$$



The image charge q'' has value,

$$q'' = -q \frac{a}{b + \delta \cos \alpha} \approx -q \frac{a}{b} + q \delta \cos \alpha \frac{a}{b^2}. \quad (3)$$

Thus, taking the limit $\delta \rightarrow 0$, $q \rightarrow \infty$ such that $q\delta = p$, the two image charges can be thought of as a dipole of strength,

$$p' = -q \frac{a}{b} \delta \frac{a^2}{b^2} = -p \frac{a^3}{b^3}, \quad (4)$$

at angle α to the radius vector as shown above, **plus** an additional charge of magnitude,

$$q''' = p \frac{a}{b^2} \cos \alpha, \quad (5)$$

located at distance a^2/b from the center. If $\alpha = \pm\pi/2$, the image of the dipole \mathbf{p} is a pure dipole.

To calculate the electric field at the center, we recall that the field of a point dipole \mathbf{p} is,

$$\mathbf{E} = \frac{3(\mathbf{p} \cdot \hat{\mathbf{r}})\hat{\mathbf{r}} - \mathbf{p}}{r^3}, \quad (6)$$

where \mathbf{r} points from the dipole to the observer. In the present problem, we define $\hat{\mathbf{z}}$ to point from the center to the dipoles, so $\hat{\mathbf{r}} = -\hat{\mathbf{z}}$. Dipole \mathbf{p} is taken to lie in the x - z plane, so that,

$$\mathbf{p} = p\hat{\mathbf{z}} \cos \alpha + p\hat{\mathbf{x}} \sin \alpha, \quad (7)$$

and,

$$\mathbf{p}' = p \frac{a^3}{b^3} \hat{\mathbf{z}} \cos \alpha - p \frac{a^3}{b^3} \hat{\mathbf{x}} \sin \alpha. \quad (8)$$

Then, the field at the center of the cavity due to the dipole, and its image dipole + extra charge q''' is,

$$\begin{aligned} \mathbf{E}(0) &= \frac{3(\mathbf{p} \cdot \hat{\mathbf{z}})\hat{\mathbf{z}} - \mathbf{p}}{b^3} + \frac{3(\mathbf{p}' \cdot \hat{\mathbf{z}})\hat{\mathbf{z}} - \mathbf{p}'}{(a^2/b)^3} - \frac{q'''\hat{\mathbf{z}}}{(a^2/b)^2} \\ &= \frac{2p\hat{\mathbf{z}} \cos \alpha - p\hat{\mathbf{x}} \sin \alpha}{b^3} + \frac{2p\hat{\mathbf{z}} \cos \alpha + p\hat{\mathbf{x}} \sin \alpha}{a^3} - \frac{p\hat{\mathbf{z}} \cos \alpha}{a^3} \\ &= p\hat{\mathbf{z}} \cos \alpha \left(\frac{2}{b^3} + \frac{1}{a^3} \right) - p\hat{\mathbf{x}} \sin \alpha \left(\frac{1}{b^3} - \frac{1}{a^3} \right). \end{aligned} \quad (9)$$