

Prospects for a Muon Collider

Kirk T. McDonald

Princeton U.

mcdonald@puphep.princeton.edu

July 17, 1998

BNL AGS/RHIC User's Meeting

Muon Collider main page:

http://www.cap.bnl.gov/mumu/mu_home_page.html

Muon Collider R&D Status Report:

http://www.cap.bnl.gov/mumu/status_report.html

Princeton muon collider page:

http://puhep1.princeton.edu/mumu/

Future Frontier Facilities

(Will the U.S. have one?)

- Hadron collider (LHC, SSC): $\approx$ \$100k/m [magnets].

  $\approx$ 2 km per TeV of CM energy.

  Ex: LHC has 14-TeV CM energy, 27 km ring, $\approx$ \$3B.

- Linear e^+e^- collider (SLAC, NLC(?)): $\approx$ \$200k/m [rf].

  $\approx$ 20 km per TeV of CM energy;

  But a lepton collider needs only $\approx 1/10$ the CM energy

  to have equivalent physics reach to a hadron collider.

  Ex: NLC, 1.5-TeV CM energy, 30 km long, $\approx$ \$6B (?).

- Muon collider: $\approx$ \$1B for source/cooler + \$100k/m for rings

  Well-defined leptonic initial state.

  $m_\mu/m_e \approx 200 \Rightarrow$ Little beam radiation.

  $\Rightarrow$ Can use storage rings.

  $\Rightarrow$ Smaller footprint.

  Technology: closer to hadron colliders.

  $\approx$ 6 km of ring per TeV of CM energy.

  Ex: 3-TeV muon collider $\approx$ \$3B (?).

Ingredients of a Muon Collider

An accelerator complex in which

- Muons (both μ^+ and μ^-) are collected from pion decay following a pN interaction.
- Muon phase volume is reduced by 10^6 by ionization cooling.
- The cooled muons are accelerated and then stored in a ring.
- $\mu^+\mu^-$ collisions are observed over the useful muon life of ≈ 1000 turns at any energy.
- Intense neutrino beams and spallation neutron beams are available as byproducts.

Muons decay: $\mu \to e\nu \qquad \Rightarrow$

- Must cool muons quickly (stochastic cooling won't do).
- Detector backgrounds at LHC level.
- Potential personnel hazard from ν interactions.

Footprints



A First Muon Collider to study light-Higgs production:



The Case for a Muon Collider

- More affordable than an e^+e^- collider at the TeV (LHC) scale.
- More affordable than either a hadron or an e^+e^- collider for (effective) energies beyond the LHC.
- Precision initial state superior even to e^+e^-.



- Initial machine could produce light Higgs via s-channel:

  Higgs coupling to μ is $(m_\mu/m_e)^2 \approx 40,000\times$ that to e.

  Beam energy resolution at a muon collider $< 10^{-5}$,

  $\Rightarrow$ Measure Higgs width.

  Add rings to 3 TeV later.

- Neutrino beams from μ decay about 10^4 hotter than present.

HEPAP Subpanel Report on PLANNING FOR THE FUTURE OF U.S. HIGH-ENERGY PHYSICS

February 1998

Recommendation on R&D for a Muon Collider

The Subpanel recommends that an expanded program of R&D be carried out on a muon collider, involving both simulation and experiments. This R&D program should have central project management, involve both laboratory and university groups, and have the aim of resolving the question of whether this machine is feasible to build and operate for exploring the high-energy frontier. The scale and progress of this R&D program should be subject to additional review in about two years.

CERN-EP/98-03
CERN-SL 98-004 (AP)
CERN-TH/98-33

Options for Future Colliders at CERN

J. Ellis, E. Keil, G. Rolandi

January 23, 1998

6 RECOMMENDATIONS

3. CERN should launch technical studies of $\mu^+\mu^-$ colliders, notably in the areas of the source and beam cooling, and should explore the possibility of locating such machines on or in the neighbourhood of the CERN site.

6. These studies should be carried out in collaborations with other laboratories, since most technical problems do not depend on the site. CERN's goal in these collaborations should be to contribute to the global pool of technologies for future collider options. It should confirm its reputation as a valuable and reliable partner in the international collaborations that will form to develop proposals for future collider projects.

The Muon Collider Collaboration

Charles M. Ankenbrandt[1], Giorgio Apollinari[2], Muzaffer Atac[1], Bruno Autin[3], Valeri I. Balbekov[1], Vernon D. Barger[4], Odette Benary[5], Scott Berg[6], Michael S. Berger[6], S. Alex Bogacz[7], T. Bolton[8], Shlomo Caspi[9], Christine Celata[9], Yong-Chul Chae[10], David B. Cline[11], John Corlett[9], Lucien Cremaldi[12], H. Thomas Diehl[1], Alexandr Drozhdin[1], Richard C. Fernow[13], David A. Finley[1], Yasuo Fukui[14], Miguel A. Furman[9], Tony Gabriel[15], Juan C. Gallardo[13] Alper A. Garren[11], Stephen H. Geer[1], Ilya F. Ginzburg[16], Michael A. Green[9], John F. Gunion[17], Ramesh Gupta[9], Tao Han[17], Katherine C. Harkay[10], Colin Johnson[3], Carol Johnstone[1], Stephen A. Kahn[13], Bruce J. King[13], Harold G. Kirk[13], Masayukiu Kumada[18], Yoshitaka Kuno[14], Paul LeBrun[1], Kevin Lee[11], Derun Li[9], David Lissauer[13], Laurence S. Littenberg[13], Changguo Lu[19], Alfredo Luccio[13], Kirk T. McDonald[19], Alfred D. McInturff[9], Frederick E. Mills[1], Nikolai V. Mokhov[1], Alfred Moretti[1], David V. Neuffer[1], King-Yuen Ng[1], Robert J. Noble[1], James H. Norem[10,1], Blaine E. Norum[20], Hiromi Okamoto[21], Yasar Onel[22], Robert B. Palmer[13], Zohreh Parsa[13], Jack M. Peterson[9], Yuriy Pischalnikov[11], Milorad Popovic[1], Eric J. Prebys[19], Zubao Qian[1], Rajendran Raja[1], Pavel Rehak[13], Thomas Roser[13], Robert Rossmanith[23], Jack Sandweiss[24], Ronald M. Scanlan[9], Lindsay Schachinger[9], Andrew M. Sessler[9], Quan-Sheng Shu[7], Gregory I. Silvestrov[25], Alexandr N. Skrinsky[25], Panagiotis Spentzouris[1], Ray Stefanski[1], Sergei Striganov[1], Iuliu Stumer[13], Don Summers[12], Valery Tayursky[25], Valeri Tcherniatine[13], Lee C. Teng[10], Alvin V. Tollestrup[1], Yağmur Torun[13,26], Dejan Trbojevic[13], William C. Turner[9], Andy Van Ginneken[1], Tatiana A. Vsevolozhskaya[25], Masayoshi Wake[14], Weishi Wan[1], Haipeng Wang[13], Robert Weggel[13], Erich H. Willen[13], David R. Winn[27], Jonathan S. Wurtele[28], Yongxiang Zhao[13], Max Zolotorev[9]

[1]*Fermi National Laboratory, P. O. Box 500, Batavia, IL 60510*
[2]*Rockefeller University, New York, NY 10021*
[3]*CERN, 1211 Geneva 23, Switzerland*
[4]*Department of Physics, University of Wisconsin, Madison, WI 53706*
[5]*Tel-Aviv University, Ramat-Aviv, Tel-Aviv 69978, Israel*
[6]*Physics Department, Indiana University, Bloomington, IN 47405*
[7]*Jefferson Laboratory, 12000 Jefferson Ave., Newport News, VA 23606*
[8]*Kansas State University, Manhattan, KS 66502-2601*
[9]*Lawrence Berkeley National Laboratory, 1 Cyclotron Rd., Berkeley, CA 94720*
[10]*Argonne National Laboratory, Argonne, IL 60439*
[11]*University of California Los Angeles, Los Angeles, CA 90095*
[12]*University of Mississippi, Oxford, MS 38677*
[13]*Brookhaven National Laboratory, Upton, NY 11973*
[14]*KEK High Energy Accelerator Research Organization, 1-1 Oho, Tsukuba 305, Japan*
[15]*Oak Ridge National Laboratory, Oak Ridge, TN 37831*
[16]*Institute of Mathematics, Prosp. ac. Koptyug 4, 630090 Novosibirsk, Russia*
[17]*Physics Department, University of California, Davis, CA 95616*
[18]*National Institute of Radiological Sciences, 4-9-1 Anagawa, Inage, Chiba, Japan*
[19]*Joseph Henry Laboratories, Princeton University, Princeton, NJ 08544*
[20]*University of Virginia, 205 McCormick Road, Charlottesville, VA 22901*
[21]*N.S.R.F, Institute for Chemical Research, Kyoto University, Gokanoshou, Uji, Kyoto 611, Japan*
[22]*Physics Department, Van Allen Hall, University of Iowa, Iowa City, IA 52242*
[23]*DESY, Hamburg, Germany*
[24]*Physics Department, Yale University, CT 06520*
[25] *Budker Institute of Nuclear Physics, 630090 Novosibirsk, Russia*
[26]*Department of Physics and Astronomy, SUNY, Stony Brook, NY 11790*
[27]*Fairfield University, Fairfield, CT 06430*
[28]*University of California Berkeley, Berkeley, CA 94720*

Spokesperson: R.B. Palmer

Meetings and Workshops

Subject	Organizer	Place	Date	Additional Information
Targetry	H. Kirk and K. McDonald	BNL	Aug. 3, 1998	Contact K. McDonald (mcdonald@puphep.princeton.eduH.)/ Kirk (hkirk@bnl.gov)
Acceleration	S. Berg and D. Neuffer	Bloomington, IN	Aug. 10-11, 1998	Contact S. Berg (jsberg@indiana.edu)
Neutrino Physics	B. King	BNL	Aug. 13-14 1998	Contact B. King (bking@bnl.gov)
Cooling Theory & Expt	R. Fernow and S. Geer	BNL	Aug. 31-Sep. 2, 1998	Contact R. Fernow (fernow@bnl.gov) or S. Geer (sgeer@fnal.gov)
Collaboration Meeting	A. Sessler	LBNL	Oct. 8-13, 1998	Contact A. Sessler (amsessler@lbl.gov)
Cooling Theory & Expt	R. Fernow and S. Geer	FNAL	Dec. 7 - 9, 1998	Contact R. Fernow (fernow@bnl.gov) or S. Geer (sgeer@fnal.gov)
Collaboration Meeting	B. Palmer	BNL	Tentative May 19-26, 1999	Contact J. Gallardo (gallardo@bnl.gov)

- PROSPECTIVE STUDY OF MUON COLLIDERS

  Introductory meeting
  Monday, July 20 1998 at 14 hrs
  CERN PS Auditorium, Bdg 6-2-024

Technical Challenges

- 16-GeV proton driver, 15 Hz, 4-MW beam power, 1-ns bunch length (C. Ankenbrandt, T. Roser...).
- **Targetry and Capture**
- **Muon Cooling**
- Acceleration – more work needed (S. Berg...)
- Storage rings have beautiful, highly corrected solutions due to heroic work of Al Garren, Carol Johnstone and Dan Trbojevič.
- Interaction region and detector design – more work needed (I. Stumer...)

A muon's view of the interaction region:



Overview of Targetry for a Muon Collider



- 1.2×10^{14} $\mu^{\pm}$/s via π-decay from a 4-MW proton beam.
- Cooling jacket around stationary target would absorb too many pions.
- Liquid-metal jet target: Ga, Hg, or solder (Bi/In/Pb/Sn).
- 20-T capture solenoid followed by a 1.25-T π-decay channel with phase-rotation via rf (to compress energy of the muon bunch).

Targetry Issues

- 1-ns beam pulse ⇒ shock heating of target.
  - Resulting pressure wave may disperse liquid (or crack solid).
  - Damage to target chamber walls?
  - Magnetic field will damp effects of pressure wave.
- Eddy currents arise as metal jet enters the capture magnet.
  - Jet is retarded and distorted, possibly dispersed.
  - Hg jet studied at CERN, but not in beam or magnetic field:



High-speed photographs of mercury jet target for CERN-PS-AA. (laboratory test)
4,000 frames per second, Jet speed: 20 ms^{-1}, diameter: 3 mm, Reynold's Number: >100,000
A. Poncet

- Targetry area also contains beam dump.
  - Need 4 MW of cooling.
  - Harsh radiation environment for magnets and rf.

An R&D Program for Targetry

at a Muon Collider

A Proposal to the BNL AGS Division

James Alessi,[b] John Corlett,[e] Richard C. Fernow,[b] Yasuo Fukui,[d] Tony Gabriel,[f] Juan C. Gallardo,[b] Michael A. Green,[e] Jerry Hastings,[b] Ahmed Hassanein,[a] Colin Johnson,[c] Stephen A. Kahn,[b] Bruce J. King,[b] Harold G. Kirk,[b] Paul LeBrun,[d] Vincent LoDestro,[b] Changguo Lu,[g] Kirk T. McDonald,[g,1] Nikolai V. Mokhov,[d] Robert B. Palmer,[b] Eric J. Prebys,[g] Claude Reed,[a] Thomas Roser,[b] Ronald M. Scanlan,[e] Dale Smith,[a] Yağmur Torun,[b,h] Andy van Ginneken,[d] Haipeng Wang,[b] Robert Weggel,[b] Yongxiang Zhao[b]

[a] *Argonne National Laboratory, Argonne, IL 60439*
[b] *Brookhaven National Laboratory, Upton, NY 11973*
[c] *CERN, 1211 Geneva 23, Switzerland*
[d] *Fermi National Laboratory, P. O. Box 500, Batavia, IL 60510*
[e] *Lawrence Berkeley National Laboratory, 1 Cyclotron Rd., Berkeley, CA 94720*
[f] *Oak Ridge National Laboratory, Oak Ridge, TN 37831*
[g] *Joseph Henry Laboratories, Princeton University, Princeton, NJ 08544*
[h] *Department of Physics and Astronomy, SUNY, Stony Brook, NY 11790*

To be submitted Sept. 1, 1998.

Studies to be performed in the AGS F.E.B. U-line.



[1] Spokesperson. Email: mcdonald@puphep.princeton.edu

Ionization Cooling

- Ionization: takes momentum away.
- RF acceleration: puts momentum back along z axis.
- $\Rightarrow$ Transverse cooling.



- Use channel of LH_2 absorbers, rf cavities and alternating solenoids (to avoid buildup of angular momentum).



- But the **energy spread rises**.
- $\Rightarrow$ Must exchange longitudinal and transverse emittance frequently to avoid beam loss due to bunch spreading.
- Can reduce energy spread by a wedge absorber at a momentum dispersion point:



- Emittance exchange via wedges + bent solenoids:



April 15, 1998

PROPOSAL

Ionization Cooling Research and Development Program for a High Luminosity Muon Collider

Charles M. Ankenbrandt[a], Muzaffer Atac[a], Giorgio Apollinari[b], Valeri I. Balbekov[a], Morris Binkley[a], S. Alex Bogacz[c], Christine Celata[d], David B. Cline[e], John Corlett[d], Lucien M. Cremaldi[f], Richard C. Fernow[g], David Finley[a], Yasuo Fukui[h], Juan C. Gallardo[g], Stephen H. Geer[a,†], Gail G. Hanson[o], Ahmed Hassanein[i], Carol Johnstone[a], Stephen A. Kahn[g], Bruce J. King[g], Harold G. Kirk[g], Thomas R. Kobilarcik[a], Yoshitaka Kuno[h], Paul LeBrun[a], Kevin Lee[e], Derun Li[d], Changguo Lu[j], Kirk T. McDonald[j], Alfred D. McInturff[d], Frederick E. Mills[a], Nikolai V. Mokhov[a], Alfred Moretti[a], Yoshiharu Mori[h], David V. Neuffer[a], Robert J. Noble[a], James H. Norem[a,i], Stephen C. O'Day[a], Yasar Onel[k], Robert B. Palmer[g], Zohreh Parsa[g], Yuriy Pischalnikov[e], Milorad Popovic[a], Eric J. Prebys[j], Zubao Qian[a], Rajendran Raja[a], Claude Reed[i], Pavel Rehak[g], Andrew M. Sessler[d], Gregory I. Silvestrov[l], Alexandr N. Skrinsky[l], Dale Smith[i], Panagiotis G. Spentzouris[a], Ray Stefanski[a], Sergei Striganov[a], Donald J. Summers[f], Lee C. Teng[i], Alvin V. Tollestrup[a], William C. Turner[d], Andreas Van Ginneken[a], Tatiana A. Vsevolozhskaya[l], David R. Winn[m], Jonathan S. Wurtele[n], Takeichiro Yokoi[h], Yongxiang Zhao[g], Max Zolotorev[d]

The MUCOOL Collaboration

[†] Spokesperson

Cooling Demonstration Experiment

Test basic cooling components:

- Alternating solenoid lattice, RF cavities, LH_2 absorber
- Lithium lens (for final cooling).
- Dispersion + wedge absorbers to exchange longitudinal and transverse phase space.

Track individual muons; simulate a bunch in software.

Possible site: Meson Lab at Fermilab:



Summary

- A muon collider offers the prospect of a more cost-effective technology for high-energy accelerators.
- Cooling the beams is the key.
- The concepts of a muon collider are still in a formative stage.
- $\Rightarrow$ Join us in exploring the physics opportunities and solving the technical challenges of a muon collider!