

EUROPEAN ORGANIZATION FOR NUCLEAR RESEARCH

CERN AT/95-06 (DI)
CERN TLS-CFM/95-06

Explosion Risks in Cryogenic Liquids Exposed to Ionising Radiation

C. R Gregory, C. W. Nuttall

Abstract

Explosions in cryogenic fluids were first reported in the early 1950's. Numerous papers have been presented describing these explosions and proposing mechanisms as to their cause. The majority of these incidents have occurred in dewars and cryogenic systems containing liquid nitrogen which have been exposed to ionising radiation.

It is now widely accepted that the explosions are caused by the very rapid decomposition of ozone, which is formed by the action of ionising radiation on oxygen dissolved in liquid nitrogen.

There is also evidence that oxides of nitrogen are formed and although it is not suggested that these compounds are the primary cause of explosions they do seem to play a catalytic role in the formation of ozone, as indeed they do in the formation of ozone in atmospheric reactions.

This paper is aimed at drawing the attention of designers of cryogenic systems in the LHC areas to the problem, in order that precautions can be taken at the design stage to reduce, or eliminate, the dangers of such explosions.

**Geneva, Switzerland
28 March, 1995**

CONTENTS

1. Introduction
2. Historic
3. Explosions
4. Physical Properties of Cryogenic Fluids
 - 4.1. General
 - 4.2. Critical Explosive Concentrations
5. Ozone formation
6. Cryostat Construction
7. Removal of ozone
8. Discussion
9. Acknowledgements
10. References

1. INTRODUCTION

Cryogenic fluids such as *Liquid Nitrogen*, *Argon* and *Helium* have been used for many years to provide low temperature environments for experimental and other purposes because they are inert, non flammable, non toxic and generally regarded as posing little hazard.

However under certain conditions explosions ^(1, 2, 3) have occurred, including some at CERN, notably in liquid nitrogen systems, after exposure to ionising radiation, for which, at the time, there was no explanation. Explosions have occurred in cryostats that have been subjected to doses of gamma radiation in the order of 10^4 Gy⁽⁴⁾, neutron fluxes of 10^{12} n.cm⁻² sec⁻¹, for a matter of two to three hours⁽⁵⁾, and irradiation by a beam of 20 μ A of 2.0 MeV electrons for 4 minutes⁽³⁾.

Radiation levels in the proposed LHC experimental areas will be much higher than those reached in existing LEP areas with projected integrated doses in the order of $2,3 \cdot 10^5$ Gy.yr⁻¹, and neutron fluxes of 10^9 n.cm⁻².sec⁻¹ ⁽⁶⁾. These values give cause for concern, particularly in the case of gamma radiation, as they are of the same magnitude as those reported in the literature where explosions have occurred.

The mechanisms of these explosions are very complex and are not yet fully understood. In this paper we review the existing literature on the subject and suggest precautions that may be required, or, further work that may be necessary to minimise explosion hazards in the LHC. Other explosions in liquefied gas containers have been caused by overpressure coupled with insufficient pressure relief, "roll-over" of the contents, or superheating resulting in the eventual catastrophic boiling of the contents. It is not intended in this paper to treat these phenomena as they are common to all cryogenic systems and are accounted for (or should be) in the design of all CERN systems.

2. HISTORIC

It might be supposed that chemical reactivity ceases at temperatures below about 100 K, but, on the contrary, many significant chemical syntheses have been described in which some operation requiring cryogenic temperatures forms a vital part of the procedure ⁽⁷⁾.

The study of chemical reactivity at cryogenic temperatures is not new, the first experiments being reported at the end of the last century, when these fluids became available in sufficiently large quantities.

Large scale use of cryogenic fluids began in the early 1950's with the advent of the space and nuclear industries, their low cost and widespread

availability has seen a multitude of uses in many fields.

3. EXPLOSIONS

As more research work was carried out, during the 1950's, into the effects of radiation on materials at cryogenic temperatures, explosions were reported for which there were no apparent explanations^(3,5,8,9). At first it was thought that the cryostat exhausts had become blocked by the formation of ice or as a result of design or operational errors, but it gradually became clear that some form of radiochemical reaction was taking place.

Liquid nitrogen was mainly used as the cooling media for these experiments, and therefore attention became focused on the possible contaminants. Commercial liquid nitrogen when delivered is usually relatively pure, however, it may easily become contaminated if it is allowed to come into contact with air. Due to their higher boiling points, oxygen and water vapour from the air are condensed and become dissolved in the liquid nitrogen.

It soon became clear that significant quantities of ozone (O_3) were being produced during the irradiation of material samples and that the ozone generation was due to the action of ionising radiation on the relatively small quantities of dissolved oxygen present in the liquid nitrogen.

The generation of ozone during the operation of an experiment is of special interest not only because of the relatively high yields of ozone which are produced when oxygen is subjected to irradiation, but in particular due to the toxic and explosive properties of the gas.

As well as the production of ozone the irradiation of liquid nitrogen may also result in the formation of oxides of nitrogen (N_2O , NO , NO_2 , N_2O_4 and N_2O_5)^(10,11) and indeed the presence of a "sludge" of oxides of nitrogen has been reported at the bottom of liquid nitrogen cryostats after irradiation⁽⁴⁾. It is considered that the oxides of nitrogen are not the cause of the explosions but they seem to enhance the yield of ozone by a catalytic effect.

It has been suggested that "active" nitrogen, consisting of atoms and excited molecules of nitrogen, may be formed and be responsible for the explosions, but this is thought unlikely as it is so reactive that the molecules will not accumulate.

Violent explosions have occurred in experiments being conducted in cryostats of liquid nitrogen, and also in small, open glass vessels containing only milligram quantities of ozone. Sufficient explosive power is generated to shatter the internal components of cryostats or cryogenic containers. It

appears that shock, local heating or even the presence of solid organic resins such as epoxies can trigger this type of explosion.

4. PHYSICAL PROPERTIES OF CRYOGENIC FLUIDS

4.1. General

The physical properties of the principal inert cryogenic fluids, together with those of oxygen, ozone and the principal oxides of nitrogen are given in Table 1.

FLUID	Boiling Pt. K (@1 atm.)	Melting Pt. K (@1 atm.)
Nitrogen (N ₂)	77.35	63
Argon (Ar)	87.29	83.2
Helium (He)	4.22	-
Oxygen (O ₂)	90.18	54.75
Ozone (O ₃)	161.3	80.7
Nitrous Oxide (N ₂ O)	182.3	170.7
Nitric Oxide (NO)	122	112
Nitrogen Dioxide (NO ₂ /N ₂ O ₄)	294.3	263.7
Nitrogen Pentoxide (N ₂ O ₅)	320	303

Table 1, Boiling and melting points of liquefied gases

Since liquid oxygen is less volatile than either liquid nitrogen or argon, oxygen enrichment occurs as the liquid nitrogen boils off. If the cryostat is refilled before all the nitrogen has been allowed to boil off, then any condensed oxygen will remain. This is even more pronounced in the case of ozone. Therefore any ozone absorbed in either liquid nitrogen or argon will remain in solution as a dissolved solid.

Liquid oxygen and ozone have a pronounced blue colour and this colour has been reported by many workers in liquid nitrogen after irradiation.


6

At ambient temperature ozone is known to be unstable, it decomposes spontaneously if only relatively slowly. However the rate of decomposition is increased by the catalytic action of certain metals, the presence of other gases

(e.g. NO), or by ionising radiation.

At temperatures below 90 K ozone is stable and does not decompose spontaneously, but under certain conditions it can decompose with explosive violence.

Explosions occur as ozone is rapidly reduced to oxygen (at a lower energy state) i.e. forming oxygen and releasing a large quantity of energy⁽¹²⁾.



$$\Delta H = - 3 \text{ kJ.g}^{-1}$$

This energy of explosion, or decomposition, is similar to the experimental value for the energy of explosion of TNT which has been measured as 4.686 kJ.g⁻¹ ⁽¹³⁾. Also the speed of detonation of pure ozone and TNT are very similar, at about 6000 m.s⁻¹.

4. 2. Critical Explosive Concentrations

Chen et al.⁽⁵⁾ postulate that the concentration of ozone needs to reach some critical value before explosions can take place. Cook et al.⁽¹⁴⁾ state that phase separation occurs giving a layer rich in ozone which enhances the probability of explosion.

Streng⁽¹⁵⁾ found that this phase separation is dependent not only upon the concentration, but also upon the temperature of the mixture of O₂/O₃, N₂/O₃ or Ar/O₃. Experimental work carried out by the French "Commissariat a l'energie atomique (CEA)" ⁽¹⁶⁾ confirms this earlier work, but explosions have been observed in Ar/O₃ mixtures at ozone concentrations below the values proposed by Streng.

	Temperature K	Theoretical min. O ₃ concentration Mole % (Streng)	O ₃ Concentration Mole % at which explosion occurred (CEA)
Oxygen	78.4	6.3	6.4
Nitrogen	77.0	4.4	5.8
Argon	85.0	8.3	4.9

Table 2, Ozone concentrations required to cause explosion

5. OZONE FORMATION

The amount of ozone formed in liquid nitrogen - oxygen mixtures exposed to ionising radiation was measured by Riley⁽¹⁷⁾. and later by

Gault⁽¹⁸⁾. It was found that $G(e)$, the number of ozone molecules formed per 100 eV, could be expressed as

$$G(e) = G_{oe} + (8,05 + 1,29 \log e)$$

This is valid for oxygen concentrations in the range 52 - 8,75. 10⁴ ppm, where G_o is the ozone yield, 12.5 molecules/100eV, from pure oxygen at 77 K, and shows that ozone production is relatively efficient even for low oxygen concentrations. For example, ozone formation in liquid nitrogen containing 10⁴ ppm oxygen, 5.7 molecules/ 100eV, is only 2.2 times that found in nitrogen containing 52ppm oxygen, 2.6 molecules/ 100eV.

The formation of ozone in liquid argon is thought to be similar to that of liquid nitrogen, but to date little concrete work has been carried out to verify this.

Experiments by Sears J. T. et al.⁽¹⁹⁾, suggest that the formation of ozone is dependent on dose at low doses but independent of dose rate, however, the steady state concentration of ozone was dependent on dose rate. This is also suggested in a theoretical study carried out by Brereton¹². Impurities also play a complex role and can radically affect the steady-state yield of ozone.

Table 3, gives the concentrations of ozone and oxides of nitrogen formed in various technical and liquefied gases on irradiation as found by Dmitriev⁽²⁰⁾. The impurity concentrations (O_2 ?) of the inert gases was assumed to be 0.5%.

Irradiated medium	O ₃ concentration at E =10 ⁴ Gy. (mg.m ⁻³)	NO ₂ concentration at E =10 ⁴ Gy . (mg .m ⁻³)
Gaseous Oxygen	9.40 .10 ²	9.7
Liquid Oxygen	2.7 .10 ⁵	2.28 .10 ³
Air	4.02 .10 ²	2.90 .10 ²
Liquid Air	1.5 .10 ⁵	5 .10 ⁴
Gaseous Nitrogen	45	11
Liquid Nitrogen	1.65 .10 ³	1.17 .10 ³
Argon	1.34 .10 ²	1.05 .10 ⁻¹
Helium	8	1.3.10 ⁻²

Table 3, Ozone formation in technical & liquefied gases

It is evident from table 3 that the ozone produced when liquefied gases are irradiated is of the order 100 times more than for the same gases at

ambient temperatures and that oxides of nitrogen are formed, albeit in much smaller concentrations than ozone. Nevertheless oxides of nitrogen may play an important role in the reactions leading to the formation of ozone.

6. CRYOSTAT CONSTRUCTION

A number of reports in the literature suggest that the materials used in the construction of the cryostats play an important role in the formation of ozone and that this formation must be, at least in part, due to surface reactions.

Work carried out by Douglas J. E. et al. ⁽²¹⁾, show that the production of ozone can be dependent on the material in contact with the fluid in the cryostat.

Unsaturated organic compounds react with ozone to produce ozonides, which are subject to sudden decomposition⁽²²⁾, and as araldite and polystyrene, which both contain double bonds, are known to play important catalytic roles in the decomposition of ozone⁽⁴⁾, this may be due to the formation of ozonides. Careful consideration should be given to their employment, and that of similar compounds, in cryostats.

Ozone may concentrate at the interface of the liquid surface and the vessel wall due to the preferential evaporation of the liquid nitrogen. It is important to choose materials with low wetting properties to limit this phenomena.

7. REMOVAL OF OZONE

Catalytic decomposition, or removal of ozone has been studied at ambient temperatures, but little work has been carried out into these processes at cryogenic temperatures.

Studies by d'Emel' Yanova et al ^(23,24) and Dewanckel et al. ⁽²⁵⁾, into the catalytic decomposition of ozone in solution, in either liquid oxygen or nitrogen, indicate that platinum, palladium and to a lesser extent silver and copper are efficient at reducing ozone to oxygen. Certain metal oxides may also be employed to destroy ozone.

Activated carbon has a high adsorptive capacity for ozone even from dilute solutions, but this should not be considered as there is a risk of accumulating relatively large quantities of ozone, with the consequent risk of explosion when warming up the system.

8. DISCUSSION

It has been shown at CERN, and at other places, that explosions in cryogenic liquids can occur when these are exposed to ionising radiation. This radiation may be gamma rays, electrons or neutrons. The cryogenic liquid in which most explosions have taken place has been liquid nitrogen, with a few in liquid argon. The mechanism for the explosions is almost certainly the

explosive decomposition of ozone, initiated by shock, or by the presence of polymeric materials in the construction of the cryostat or the presence of foreign matter such as dust particles acting as catalysts. The ozone is formed by the irradiation of oxygen impurities dissolved in the cryogenic liquid. There is some evidence that nitrogen plays a role in the reactions leading to the formation of ozone, but the preponderance of explosions in liquid nitrogen systems may simply be due to the fact that many more liquid nitrogen systems have been exposed to ionising radiation than have, for instance, liquid argon systems.

The quantity of ozone formed is a function of the total dose and very much less a function of the initial oxygen content. The effect of the dose rate is less clearly understood.

The above mechanism appears to represent a serious risk to the cryogenic systems being proposed for the LHC experiments, since quantities of cryogenic liquids will be found in zones of high radiation (e.g. EM Calorimeter end caps). The total annual dose will fall within the range known to create dangerous amounts of ozone in liquid nitrogen, or, argon and the complexity of the system could well facilitate the contamination of the cryogens by oxygen unless special precautions are taken.

It is of primary importance that these problems be addressed at the design stage, to reduce the potential danger by minimising the quantity of oxygen within the radiation field, for example using closed cycle secondary circuits filled with pure liquids cooled by heat exchange with commercial liquid nitrogen not exposed to ionising radiation.

It would be wise to monitor continuously the oxygen level in the cryogens exposed to ionising radiation and it may be necessary to design in systems for the removal of ozone as it is formed should the studies show that the risk to the cryogenic systems in the LHC experiments is such that it cannot be accepted.

This report has concentrated on the experimental areas where liquid nitrogen and argon will be present, for it is in these systems that most explosions have occurred. As far as we are aware there have been no reported incidents due to the problem of ozone formation and explosive decomposition in irradiated liquid helium. However there have been reports of ozone formation and explosion in solid air condensed on external cold spots of liquid helium cryogenic systems exposed to ionising radiation⁽²⁷⁾. This

10

problem should be taken into account in the design and operation of the cryogenic systems of the LHC machine.

As stated above, the total dose plays an important role in the formation of ozone, although the effects of dose rate and oxygen concentration are less clearly defined. The role played by nitrogen oxides in the formation of ozone

is also unclear, as although these compounds do not appear to be the source of explosions, they may catalyse the formation of ozone. Further experimental work into the effects of dose rates and oxygen concentration would be of great value, as would work on the risks in liquid argon systems as much less information exists on this latter subject.

9. ACKNOWLEDGEMENTS

We are indebted to S. Brereton, of Lawrence Livermore Laboratory, L. C. Cadwallader of Idaho National Engineering Laboratory and G. Bon Mardion, of Commissariat a l'energie atomique, service des basses temperatures, for their contributions with respect to bibliographic references. These have enabled us to do a more complete search than we would otherwise have been able to do. Also T. Niinikoski, L. Leistam and K. Potter for their help and encouragement.

10. REFERENCES

1. Niinikoski T. CERN. Memo to C. W. Nuttall, 17th- June 1994.
2. Bochirol L. et al.. A cryogenic device for irradiation in liquid nitrogen, Cryogenics, September 1960.
3. Loomis J. S. et al. Ozone hazards incurred by electron van de GRAAFF

operation, Nuc. Inst. & Methods, 15,1962.

4. Walker P. T. Explosions in irradiated liquid nitrogen, LJKAEA, research group, Harwell, January 1961.
5. Chen C. W. et al. On the cause of explosions in reactor cryostats for liquid nitrogen, Cryogenics, April 1969.
6. ATLAS Technical Proposal, CERN/LHCC/94-43, 15th December 1994.
7. McGee H. A. et al. Cryochemistry, Cryogenics, September 1962.
8. Cottrell W. B. ORNL Report CF 58-7-97(1958).
9. Heyne L. The concentration of oxygen in nitrogen cryostats for reactor irradiations, Cryogenics, December 1962.
10. Junk G. et al, The presence of N³⁺ and N⁴⁺ in the mass spectra of molecular nitrogen. J. of Am. Chem. Soc. Vol 80, p. 2908,1958.
11. Kelly R et al, The kinetics of the decay of nitrogen atoms as determined from chemical measurements of atom concentrations as a function of pressure. Can. J. Chem. Vol. 37, p. 62,1959.
12. Brereton S. J. Explosion hazard in liquid nitrogen cooled fusion systems, Paper, Idaho National Engineering Lab.
13. F. P. Lees, Loss Prevention in the process Industries Butterworths, London, 1983.
14. Cook G. A. et al. Explosion limits of oxone- oxygen mixtures, Ind. & Eng. Chem. 48, 4,736-741, April 1956.
15. Streng A. G. Tables of ozone properties. J. of Chem. and Engineering data, 6, No 3, 431 - 436, July 1961.
16. Dewanckel B. Explosibilite de l'ozone en solution dans l'oxygene, l'azote et l'argon liquides. Centres d'etudes nucleaires de Grenoble. Service des basses temperatures, Note SBT/CT/82 - 95. Mars 1982.
17. Riley J. F. Radiolysis of liquid oxygen and oxygen - nitrogen at 770K, USAEC Report ORNL - 3488, 42 - 43.1963.
18. Gault J. D. et al, Ozone formation by the radiolysis of liquid nitrogen; Calculation and measurement. Nuclear safety, Vol. 14, No 5, September -October 1973.
19. Sears J. T. et al, Radiolytic formation and decomposition of ozone. J. of Physical Chem. Vol. 72, No 4, April 1968.

20. Dmitriev M. T. Formation of ozone and nitrogen oxides in technical and liquefied gases under irradiation conditions. *J. of Applied Chem. of the USSR* Vol. 41, No 1, January 1968.
21. Douglas J. E. et al, Peculiarities of ozone formation in electron irradiation of nitrogen - oxygen mixtures. *J. Chem. Phys.* p. 1416, November 1959.
22. *Encyclopedie des gaz, L'air Liquide*, Elsevier science pub B. V. Amsterdam. 1976.
23. Emel' Yanova G. I. et al, Low temperature heterogeneous catalytic decomposition of concentrated liquid ozone. *Russian J. of Phys. Chem.*, January 1964.
24. Emel' Yanova G. I. et al, Mechanism and kinetics of low temperature catalytic decomposition of liquid ozone on platinum and palladium. *Russian J. of Phys. Chem.* October 1965.
25. Dewanckel et al, Destruction catalytique de l'ozone en solution dans l'oxygene liquide. Centres d'etudes nucleaires de Grenoble. Service des basses temperatures, Note SBT/CT/81- 82, octobre 1981.
26. Niinikoski T. Memo to C. W. Nuttall, 26th- September 1994.
27. Cadwallader L. C. Cryogenic system operating experience review for fusion applications. Idaho National Engineering Laboratory, EGG-FSP-10048, January 1992.